

NARRATIVE REVIEW

THE THERAPEUTIC VALUE OF LAUGHTER IN MEDICINE

Ramon Mora-Ripoll, MD, PhD

Objective • The aim of this review is to identify, critically evaluate, and summarize the laughter literature across a number of fields related to medicine and health care to assess to what extent laughter health-related benefits are currently supported by empirical evidence.

Data Sources and Study Selection • A comprehensive laughter literature search was performed. A thorough search of the gray literature was also undertaken. A list of inclusion and exclusion criteria was identified.

Data Extraction • It was necessary to distinguish between humor and laughter to assess health-related outcomes elicited by laughter only.

Data Synthesis • Thematic analysis was applied to summarize laughter health-related outcomes, relationships, and general robustness.

Conclusions: Laughter has shown physiological, psychological,

social, spiritual, and quality-of-life benefits. Adverse effects are very limited, and laughter is practically lacking in contraindications. Therapeutic efficacy of laughter is mainly derived from spontaneous laughter (triggered by external stimuli or positive emotions) and self-induced laughter (triggered by oneself at will), both occurring with or without humor. The brain is not able to distinguish between these types; therefore, it is assumed that similar benefits may be achieved with one or the other. Although there is not enough data to demonstrate that laughter is an all-around healing agent, this review concludes that there exists sufficient evidence to suggest that laughter has some positive, quantifiable effects on certain aspects of health. In this era of evidence-based medicine, it would be appropriate for laughter to be used as a complementary/alternative medicine in the prevention and treatment of illnesses, although further well-designed research is warranted. (*Altern Ther Health Med.* 2010;16(6):56-64.)

Ramon Mora-Ripoll, MD, PhD, is medical scientific director at Organización Mundial de la Risa, Barcelona, Spain.

Corresponding author: Ramon Mora-Ripoll, MD, PhD
E-mail address: ramon.morari@gmail.com

That laughter has health benefits has been claimed for centuries; however, during the past decades, several laughter- and humor-based interventions have gained widespread acceptance, and scientific studies of this phenomenon have generated considerable medical and public interest. In 1976, Norman Cousins published the article "Anatomy of an Illness" in *The New England Journal of Medicine*, in which he explained that 10 minutes of genuine belly laughter as a result of viewing comic films had a considerable analgesic effect and gave him at least 2 hours of sleep free from the pain of ankylosing spondylitis.¹ It was never clear whether his pain relief was due to the laughter or to the massive dose of vitamin C administered simultaneously, but the case became famous and encouraged scientists to investigate and define how laughter may heal, founding the basics of current "laughter medicine."

Though it takes skill and time to develop the science and art of what makes a person laugh and to ascertain the related benefits, laughter therapy does not require large amounts of time or money to be applied. Unlike other therapies that are more time-

consuming or expensive, the use of laughter can be implemented easily and cost-effectively in patient care.^{2,3} Nevertheless, there are now so-called "laughter trainers" and accounts all over the popular media proclaiming the supposed physical, psychological, emotional, social, occupational, spiritual, and quality-of-life benefits of laughter. Scientific data for these claims must be well documented before evidence-based "laughter medicine" can be widely supported by the health care community.^{4,9}

The aim of this review is to identify, critically evaluate, and summarize the laughter literature across a number of fields related to health, health care, patient care, and medicine with the purpose of assessing to what extent laughter health-related benefits are currently supported by empirical evidence.

METHODS

Definitions

This review focuses on the health-related effects of laughter only. One of the most significant methodological problems in laughter research is the failure to distinguish humor from laughter. A full discussion on laughter and humor theories is beyond the scope of this review, but a few basic definitions are essential.⁶ For the purpose of this review, *humor* is defined as one of the stimuli that can help people laugh and feel happy. *Sense of humor* is a psychological trait that varies considerably and allows people to respond to different types of humorous stimuli. And *laughter*

is defined as a psychophysiological response^{10,11} to either humor or any other stimuli with the following characteristics¹²: (1) powerful contractions of the diaphragm together with repetitive vocal sounds produced by the action of the resonating chambers of the pharynx, mouth, and nasal cavities; (2) typical facial expression (motion of about 50 facial muscles, mainly around the mouth), which may include the release of tears; (3) motion of several groups of muscles of the body (more than 300 may be distinct); and (4) a sequence of associated neurophysiological processes (cardiovascular and respiratory changes, activation of neuroendocrine and immune circuits). Internally, laughter comes with positive psychological shifts and a subjective identifiable emotion (hilarity) that has been compared to the one coming from sexual activity or other joyful bodily responses.

It is apparent that humor and laughter are distinct events (although often associated). Whereas humor is a stimulus and can occur without laughter, laughter is a response and can occur without humor.¹³ It is necessary to distinguish between these variables, as many analyses of humor have used a humor stimulus (such as a comic movie) to determine the effect of "humor" on a health-related outcome, and others look specifically at the effects of laughter on these outcomes. Still others explore different ways to assess sense of humor in an attempt to analyze whether scoring higher on a sense-of-humor scale is associated with certain health outcomes.

Design

A systematic review was conducted. This type of review is particularly useful where the aggregation of data is difficult because diverse definitions, many studies or fields, and different outcomes are being analyzed.¹⁴ A comprehensive laughter and humor literature search was performed using a variety of databases and keywords. A manual search of relevant sources, a journal-specific search, and a manual search of references included in relevant retrieved articles also were included (Table 1). A thorough search of the gray literature also was undertaken (eg, Google). A list of inclusion and exclusion criteria was identified (Table 2). All relevant published articles up to 2008 were reviewed. No papers were excluded on the basis of quality because of the dearth of literature meeting the established inclusion or exclusion criteria. Thematic analysis was applied to summarize laughter and humor theories, health-related outcomes, patient outcomes, relationships, and general robustness.¹⁵ This review was not funded.

RESULTS

The Effects of Laughter on Health Outcomes

The field of medicine is surprisingly less present in laughter research than would be expected. Its research is mainly dominated by the field of (medical) psychology across psychobiological, social, and health domains.

Mechanism of Action

In pursuing the therapeutic efficacy of laughter, four poten-

TABLE 1 Databases, Journals, and Keywords Searched

Databases	Journals specific	Keywords
Cochrane Library	<i>Altern Ther Health Med</i>	Laughter (or laugh or laughing)
Medline	<i>BMC Complement</i>	
PubMed	<i>Altern Med</i>	Humor (not aqueous, not vitreous)
SAGE Journals	<i>Complement Ther</i>	
Science Direct	<i>Clin Pract</i>	Mirth (or mirthful)
Internet (Google)	<i>Complement Ther Med</i>	Therapy
	<i>Evid Based Complement</i>	(or intervention)
	<i>Alternat Med</i>	Complementary
	<i>J Altern Complement Med</i>	medicine
	<i>J Psychosom Res</i>	Alternative medicine
	<i>Int J Humor Res</i>	Holistic medicine
		Integrative medicine
		Mind-body
		Medicine
		Health (and health care)
		Patient (and patient care)

TABLE 2 Laughter Literature Search: Inclusion and Exclusion Criteria

Inclusion

- Laughter or humor intervention main focus of paper
- Direct or indirect relationship to health outcomes and medicine/therapy
- Paper either research or seminal in some form
- Literature review of >15 references (medicine texts only)

Exclusion

- Humor stimulus or intervention not eliciting or measuring laughter
- Brief items, anecdotes, cartoons, pictures, obituaries, commentaries, or interviews
- Brief literature reviews based on ≤15 references (medicine texts only)
- Semantics, linguistics, history (nontherapeutic research), eg, joke constructions, entertainment
- Textbooks, media-related items, including health promotion
- Non-health care occupation-related items
- Non-English, non-Spanish language publications

tial mechanisms of action have been established that would demonstrate its direct or indirect health benefits (humor-health connection).^{4,8} First, laughter can lead to direct physiological changes to the muscular, cardiovascular, immune, and neuroendocrine systems, which would have immediate or long-term beneficial effects to the body.^{16,17} Accordingly, laughter is crucial in this model and may be expected to have beneficial health outcomes even without humor, as advocated by the laughter club movement that originated in India in 1995.¹⁸ The more you laugh, the more benefits you obtain. Secondly, laughter can lead to more positive emotional states, which also may have direct benefits to health or contribute to a personal perception of better health or quality of life.¹⁹ Laughter is not that essential in this second model, as positive emotions may also be elicited by humor (without laughter), amusement, happiness, joy, love, and others. Here, the more playful approach to life, the more benefits. Third,

laughter can optimize one's own strategies for coping with stress and strengthen personal pain tolerance,²⁰ which may reduce the negative impact on health benefits that both can have. According to this stress-moderator model, which provides indirect effects, laughter during nonstressful times would be less relevant to health. Finally, laughter may indirectly increase one's social competencies, which as a result may increase interpersonal skills. In turn, the greater levels of social support gained may confer stress-buffering and health-enhancing benefits.^{21,22} Laughter's role is here a lot less patent, as the main focus is on social skills. No other potential mechanisms of action for laughter have been reported as yet.

Therapeutic Efficacy

Although humor and laughter have been used therapeutically in a variety of medical and other conditions, well-designed randomized controlled trials (RCTs) have not been conducted to date validating the therapeutic efficacy of laughter, and only very few trials have been performed otherwise. However, health outcomes have been reported in multiple areas of medicine and patient care. This review has identified health-related laughter research, excluding pathological laughter, in the following areas: oncology²³⁻²⁶; allergy and dermatology^{27,28}; immunology^{29,30}; pulmonology³¹⁻³³; cardiology, endocrinology, and metabolism^{17,34-36}; internal medicine and rheumatology³⁷; rehabilitation^{38,39}; psychiatry and medical psychology⁴⁰⁻⁴³; anatomy, neurology, and imaging⁴⁴⁻⁵⁰; biophysics and acoustics⁵¹⁻⁵⁵; geriatrics and aging⁵⁶⁻⁵⁸; pediatrics⁵⁹⁻⁶¹; obstetrics⁶²; surgery⁶³⁻⁶⁵; dentistry⁶⁶⁻⁶⁸; nursing⁶⁹⁻⁷⁴; critical, palliative, and terminal care⁷⁵⁻⁸²; hospice care⁸³⁻⁸⁵; home care⁸⁶; general patient care and primary care⁸⁷⁻⁹⁰; epidemiology and public health^{91,92}; complementary and alternative medicine (CAM)^{93,94}; and medical and health sciences training.⁹⁵⁻⁹⁷

Physiological Benefits of Laughter

Humor, mirth, and laughter have numerous effects involving the muscular, cardiovascular, respiratory, endocrine, immune, and central nervous systems. The effects of laughter on certain physiological outcomes are briefly summarized in Table 3. The research reviewed in this area relates to the impact of laughter on the entire body and can be lumped into the following main physiological effects: laughter (1) exercises and relaxes muscles, (2) improves respiration, (3) stimulates circulation, (4) decreases stress hormones, (5) increases the immune system's defenses, (6) elevates pain threshold and tolerance, and (7) enhances mental functioning.

Psychological Benefits of Laughter

The psychological effects of humor and laughter relate primarily to both as a coping mechanism and, to a lesser extent, their enhancement of interpersonal relationships. Table 4 provides a brief overview of the effects of laughter on particular psychological outcomes. The research reviewed in this area, somehow larger and much stronger than the evidence for the physiological health benefits, can be summarized as follows: laughter (1) reduc-

es stress, anxiety, and tension and counteracts symptoms of depression; (2) elevates mood, self-esteem, hope, energy, and vigor; (3) enhances memory, creative thinking, and problem solving; (4) improves interpersonal interaction, relationship, attraction, and closeness; (5) increases friendliness and helpfulness and builds group identity, solidarity, and cohesiveness; (6) promotes psychological well-being; (7) improves quality of life and patient care; and (8) intensifies mirth and is contagious.

Safety

The side effects of laughter are very limited. In specific cases, the appearance of a laughter-induced syncope has been reported.¹⁶⁷⁻¹⁷⁰ Contraindications are nearly nonexistent; however, precaution is advised with patients who were recently released from surgery or who have certain cardiovascular or respiratory diseases or glaucoma.

Laughter Types and Health Benefits

Several kinds of laughter have been identified depending on various parameters and different fields of the scientific research.¹⁷¹⁻¹⁷³ From a medical and therapeutic point of view, five large groups can be summarized¹²: (1) genuine or spontaneous laughter, (2) self-induced simulated laughter, (3) stimulated laughter, (4) induced laughter, and (5) pathological laughter. Spontaneous laughter, unrelated to one's own free will, is triggered by different (external) stimuli and positive emotions (ie, happiness, mirth, joy, fun, triumph, humor, surprise, emotional release, or by contagion). It has been reported that spontaneous laughter causes typical contractions of the muscles around the eye socket (Duchenne laughter/smile¹⁷⁴). Self-induced simulated laughter is triggered by oneself at will, with no specific reason (purposeful, unconditional) and therefore not elicited by humor, fun, other stimuli, or positive emotions. Stimulated laughter happens as a result of the physical (reflex) action of certain external factors (ie, to be ticklish, particular facial or bodily motions, by pressing laughter bones¹⁷⁵). More superficial and empty-headed, induced laughter is the consequence of the effects of specific drugs or substances (ie, alcohol, caffeine, amphetamines, cannabis, lysergic acid diethylamide [LSD], nitrous oxide or "laughing gas," and others). Finally, pathological laughter is secondary to injuries to the central nervous system caused by various temporary or permanent neurological diseases and also may occur with certain psychiatric disorders. Pathological laughter is developed with no specific stimulus; is not connected with emotional changes; has no voluntary control of its duration, intensity or facial expression; and sometimes comes with "pathological crying."¹⁷⁶

The therapeutic benefits of the different types of laughter concern in particular the first two, spontaneous laughter and self-induced simulated laughter, and stimulated laughter to a lesser extent. Table 5 shows the main characteristics of the first two types of laughter. Spontaneous laughter and self-induced simulated laughter are not that different than one might believe initially. The only clear difference is in the initial stage of providing a stimulus and the triggering of laughter. In the first case, an external

TABLE 3 Effects of Laughter on Health-related Physiological Outcomes*

Physiological Outcome	Intervention and Results
Muscle relaxation	Periods of intense laughter are followed by relaxed muscle tone ⁹⁸ or H-reflex depression. ⁹⁹
Heart rate, respiratory rate, blood pressure, EE, oxygen levels	Laughter led to immediate increases in heart rate, respiratory rate, and oxygen consumption ¹⁰⁰⁻¹⁰⁴ and may improve oxygen saturation levels ¹⁰⁵ ; laughter did not significantly affect heart rate ¹⁰⁶ or oxygen consumption ³⁴ ; following laughter, there is a corresponding decrease in heart rate and respiratory rate ¹⁶ ; laughing had an acute effect on systolic blood pressure ^{100,107} ; no significant effect ^{34,108} ; laughter with physical exercise may be effective to lower the blood pressure as a long-term effect ¹⁰⁹ ; genuine voiced laughter caused a 10% to 20% increase in EE and heart rate above resting values, which means that 10-15 min of laughter per day could increase total EE by 10-40 kcal. ¹¹⁰
Effect on cardiovascular performance	Laughter increased stroke volume and cardiac output, and decreased arterial-venous O ₂ difference and systemic vascular resistance ³⁴ ; laughter elicited by cinematic viewing improved endothelial-dependent vasodilation ¹¹¹ ; laughter decreased levels of serum cortisol and plasma von Willebrand factor ¹¹² ; laughter induced by a comic movie led to a significant decrease in aortic stiffness and wave reflections. ¹¹³
Cardiovascular protection (long-term effects)	Inverse association between propensity to laugh and coronary heart disease ¹¹⁴ ; fewer arrhythmias and recurrences of myocardial infarctions during cardiac rehabilitation (after myocardial infarction) when self-selected viewed humor was used as an adjunct to standard therapy ¹¹⁵ ; mirthful laughter led to lower the incidence of myocardial infarction in high-risk diabetic patients. ¹¹⁶
Endocrine stress markers (cortisol, CgA) and various hormonal measures	After watching a comic film, laughter reduced serum cortisol levels, ^{117,119} increased salivary CgA levels, ¹²⁰ increased urinary excretion of epinephrine and norepinephrine ¹²¹ ; appeared to reduce serum levels of dopac (dopamine catabolite), epinephrine, and HGH ¹¹⁸ ; no significant changes were found in serum prolactin, beta-endorphins, ACTH, and norepinephrine ^{117,118} ; laughter increased beta-endorphins and HGH ¹²² ; laughter elevated breast-milk melatonin in both healthy and atopic eczema mothers ²⁸ ; behavior of perceptual anticipation of mirthful laughter decreased serum cortisol, epinephrine, and dopac. ¹²³
Neuroimmune parameters: salivary IgA (SIgA), serum immunoglobuline levels, NK cell activity, leukocyte population	Exposure to a humorous stimulus increased NK activity ^{29,117,124} , did not significantly increase NK activity ¹²⁵ ; increased SIgA ^{30,126-129} , increased serum IgA, IgG, IgM ¹⁴⁷ ; relative increase in total leukocytes and specific leukocyte subsets. ¹¹⁷
Pain threshold and tolerance	Laughter (film-induced) increased pain tolerance and discomfort thresholds. ¹³⁰⁻¹³⁵
Effects in asthma and COPD patients	Viewing a humorous film decreased bronchial responsiveness in asthmatic patients ¹³⁶ ; laughter and smiling induced by a humor intervention (clowning) were able to reduce hyperinflation in severe and very severe COPD patients. ¹³⁷
Effects in patients with rheumatoid arthritis (neuroimmune parameters)	Mirthful laughter decreased serum proinflammatory cytokine levels, ¹³⁸ growth hormone, and IGF-1 ¹³⁹ ; increased antiinflammatory cytokine levels ¹³⁸ ; or reduced serum interleukin-6 levels. ^{37,140}
Effects in type 2 diabetes patients (blood glucose levels, neuroimmune parameters, others)	Inhibitory effect of laughter (elicited by a comedy show) on the increase of postprandial blood glucose level ¹⁴¹ ; laughter influenced the gene expression profile in the peripheral blood leukocytes ¹⁴² ; laughter may prevent the exacerbation of diabetic nephropathy ³⁶ and diabetic microvascular complications ¹⁴³ ; laughter may contribute to amelioration of postprandial blood glucose elevation through a modulation of NK cell activity caused by upregulation of relating genes ¹⁴⁴ ; mirthful laughter led to lower the serum epinephrine and norepinephrine levels, decreased inflammatory cytokines and C-reactive protein, and increased HDL cholesterol in high risk diabetic patients with hypertension and hyperlipidemia. ¹¹⁶
Effects in atopic dermatitis patients (allergy parameters)	Laughter and humor reduced allergen-induced wheal reactions, ²⁷ reduced allergen-specific IgE production, ¹⁴⁵ improved night-time wakening, ¹⁴⁶ and reduced serum neurotrophin levels. ¹⁴⁷
Other: skin response, binocular rivalry, diaphragm electromyography	Laughter increased galvanic skin response ¹⁴⁸ or conductance, ¹⁴⁹ indicating activation of sympathetic nervous system; laughter episodes (while narrating jokes) led to cessation of binocular rivalry ¹⁵⁰ ; hearty laughter (while viewing a comic video) elicited waving patterns clearly different from those of coughing or sneezing. ¹¹³

*EE indicates energy expenditure; CgA, chromogranin A; HGH, human growth hormone; ACTH, adrenocorticotropic hormone; Ig, immunoglobulin; NK, natural killer; COPD, chronic obstructive pulmonary disease; IGF-1, insulin-like growth factor 1; HDL, high-density lipoprotein.

TABLE 4 Effects of Laughter on Health-related Psychological Outcomes

Psychological Outcome	Intervention and Results
Effects on mood, stress, depression and/or anxiety symptoms	Laughter improved mood and positive affect in healthy adults ¹⁵¹⁻¹⁵³ ; temporarily improved depressed mood in depression ¹⁵⁴ ; moderated stress in healthy adults ^{112,155,156} or anxiety. ^{106,157}
Effects on psychotic symptoms	In patients with schizophrenia, a humor and laughter intervention reduced hostility and depression/anxiety scores; improved activation scores and social support ^{41,158} ; lowered the levels of psychopathology; and improved social competence. ¹⁵⁹
Performance, personal efficacy, coping abilities	Purposeful laughter significantly increased different aspects of self-efficacy, including self-regulation, optimism, positive emotions, and social identification, and maintained these gains at follow-up ¹¹² ; laughter and humor improved coping abilities. ^{40,160,161}
Psychotherapy, group therapy, desensitization	Conversational laughter helped prevent or resolve risk of confrontation in addiction group therapy ¹⁶² ; humor desensitization reduced fear as effectively as traditional techniques. ¹⁶³
Quality of life, patient care, well-being	Laughter and humor improved quality of life in depressed patients ^{164,165} ; promoted psychological well-being and enhanced patient care in different clinical settings. ^{25,39,58,78,79,166}

stimulus (coming from other than the laugher) is commonly provided, and laughter is triggered. In the second case, it occurs by the laugher him/herself (purposeful laughter). Modern laughter therapy is based on the following fundamental principle: through several exercises, techniques, activities, and dynamics, a person or a group of people is taken to a feeling of lack of inhibitions to achieve the binomial self-induced laughter–spontaneous laughter and to experience its physical, psychological, emotional, and spiritual benefits. The human brain is not able in the end to distinguish spontaneous from self-induced laughter (“motion creates emotion” theory); therefore, their corresponding health-related benefits are alleged to be alike, as some authors contend.¹⁸ However, further research is warranted to confirm this assumption (no studies were identified in this review). Indeed, self-induced stimulated laughter may lead to a higher “laughter exposure” both by achieving greater intensity and duration at will or by triggering contagious and turning into spontaneous laughter, which might create greater accompanying psychophysiological changes.^{4,16} As a laughter type, self-induced simulated laughter is becoming increasingly popular worldwide, as it is the foundation of the Laughter Club movement (Laughter Yoga).^{18,177}

Laughter Research Pitfalls and Flaws

Though some of the studies on the effects of laughter

TABLE 5 Laughter Medicine: Main Characteristics of Spontaneous Laughter and Self-induced Simulated Laughter

	Spontaneous Laughter	Self-induced Laughter
Genuine expression of positive emotions	Yes	Sometimes
Humor-related	Very often, but not always	Rarely
Laughing “at” or “with” others	Commonly “at”	Always “with”
Personal engagement/effort to think	Often necessary	Rarely necessary
Presence of “Duchenne laughter/smile”	Very often (when not fake)	Often
Together with bodily motion	Yes (uncontrolled)	Yes (controlled)
One’s own free will	No, unrelated	Yes, purposeful
Triggering off stimulus	Identifiable (external)	None (self-induced)
Contagious	Yes (often)	Yes (very often)
Self-control on intensity and duration	Minimum or less	Maximum or more
Presence of vocal sounds	Common (50%) ¹⁹⁷	At will
Most common vocalizations	Ha/ho	At will: ha/he/hi/ho/hu and others
Evidence of therapeutic value in medicine	Some studies conducted	Very few studies
Best dynamics for experimenting laughter	In a group	Alone/in a group

*Contraction of the muscles around the eye socket.

marked a new start in research called psychoneuroimmunology,¹⁷⁸ which explores the interactions between the central nervous system and the immune and endocrine systems, not all of these research attempts have been completely successful. The highest-quality studies were executed on the effects of a humorous stimulus (comic movie) and subsequent laughter on pain tolerance, which provide strong evidence of increased long-term effects not merely due to distraction.⁷ However, there is still not enough empirical evidence that hearty laughter has pain-killing effects or that laughter stimulates the production of endorphins, other neurohormones (human growth hormone, oxytocin, melatonin, prolactin, adrenocorticotrophic hormone) and neurotransmitters (serotonin, dopamine, others).^{3,9,179} The weakest investigations are those on endocrine stress markers (serum cortisol, salivary, chromogranin A levels, others) and immune system parameters.^{4,6} The results were not consistent or conclusive because methodological flaws might have prevented the expected physiological “benefits” from being detected.¹⁹

The criticisms of the weaknesses and quality of the studies conducted are focused mainly on internal design flaws and invalid/lack of generalizability of results. The main internal flaws identified, some of them reported elsewhere,^{5,8} included the following: small sample size (≤ 10 participants), no randomized design, inadequate or nonexistent control groups, no standardized baseline measurement, unreliable measures of blood and saliva assays, no statistical tests or too many, failure to distinguish laughter from humor, failure to confirm the presence of laughter, suboptimal laughter exposure (either insufficient duration or intensity), failure to differentiate short-term from long-term effects, and many other confusion factors. Collectively, current available data reviewed suffer simply from too few well-designed studies to draw valid conclusions about some of the health-enhancing changes produced from laughter.¹⁷⁹

DISCUSSION

In this review, the focus was placed on laughter research occurring either in the presence or humor or in its absence and related health benefits. This distinct approach makes this narrative review unique, as most of the previous studies and reviews have assessed mainly (sense of) humor interventions and their outcomes, where, unfortunately, laughter, as a common response to humorous stimuli, was not always present, measured, or monitored.⁴ However, this review may have been limited by the amount of humor and laughter literature available; the numerous definitions and fields and different methodologies and outcomes from where it emerged; the failure to adjust for the presence, duration, or intensity of laughter and corresponding effects; and the overall complexity of the laughter phenomenon itself.

Laughter Research Challenges and Future Directions

Laughter research designs can be quite challenging. As previous studies indicate, some individuals who are exposed to a humorous stimulus do not always laugh. To help control for this, it is crucial that future laughter research includes some measure of subject response to the humorous stimuli whenever it is used to help elicit laughter. An alternative design to help preclude laughter absence and to adjust for its intensity and duration is to conduct self-induced simulated laughter intervention trials. Although it has been reported that very few people are able to convincingly laugh on command,⁷ self-induced laughter is entirely achievable and appears to be the most realistic, sustainable, and generalizable intervention to be used in future laughter research. In this review, only two studies experimenting with purposeful laughter and then assessing its health effects were identified.^{180,181} Another methodological concern is that of control groups. The need for at least two additional control groups has been suggested⁶: a negative one to control for the effect of general emotional arousal and a positive one to control for positive emotions that are not necessarily laughter-oriented. Larger samples of healthy subjects and trials in different clinical populations also are warranted. Therefore, it may be useful for researchers in an area dominated by psychology to carry out interdisciplinary

studies involving experts from different health care fields.

Laughter researchers also may have difficulty defining precise and measurable outcomes for a therapy for which the main effect is often subjective or dependent on the skill of the practitioner and the laughter him/herself. Laughter therapy often has to be tailored for an individual's specific needs and may not be able to be studied at a conventional "active ingredient" or "dose" level. Moreover, individuals frequently use a variety of CAM modalities simultaneously or adjunctively with conventional therapies, which may attenuate or magnify treatment effects. Some of these challenges have not been properly addressed through alternative study designs. Collectively, research on laughter is still in its infancy, and many efforts are required to enhance the quality and validity of trial designs and health-related outcomes, with the imperative need of distinguishing (sense of) humor and laughter. Clearly, more groundwork is needed to determine the best methods of assessing and documenting health-related outcomes on laughter in different patient populations. One of the first steps may have been performed by Kimata et al,¹⁸² who recently published some remarkable results on "diaphragm EMG (electromyogram)," the first exact system for the measurement of laughter itself. This measuring system can specify the starting point and duration of laughter precisely of 1/3000 second; therefore, it will make detailed analysis of the healthy effects of laughter possible in different clinical settings. Furthermore, the authors state that this method is suitable as a precise way to assess any kind of laughter and does not require any special medical or mechanical techniques, so it also may be useful for other, nonmedical purposes.

Laughter as Medicine

Western thinking around laughter as a medicine began to crystallize in 1976 when Cousins published his "Anatomy of an Illness." However, inviting and facilitating laughter in therapy is not the same as developing and using humor to make the patient laugh. Humor is not necessary to have subsequent laughter. Adults can laugh without it, as do infants and children. While laughter medicine takes skill and time to be developed, laughter therapy itself can be implemented easily and is cost-effective in patient care.^{2,3} Health care professionals do not have to be stand-up comics, clowns, or magicians to bring laughter into clinical settings. Just acknowledging how important laughter can be and having a cheerful and spirited approach is a good place to start. After all, half of the benefit of laughter, in addition to healing, is sharing it.¹⁸³ Nevertheless, the health community is still slow in accepting and considering laughter as a healing tool within CAM.¹⁸⁴ In order to offer patients the benefits of laughter, health care professionals must be willing to break loose from conventional therapeutic constraints, regain their own laughter, and learn the techniques to facilitate laughter in their patients. Laughter deserves a special place in medical practice and daily life. This is the mission of the Association for Applied and Therapeutic Humor, an international community of medical, education, and public speaking professionals who study, practice,

and promote healthy humor and laughter. Based in Spain, the “Organización Mundial de la Risa” also is investing in laughter research and training.

Laughter Medicine for Health Care Professionals

Laughter is important in medicine and may enhance conversation between health care professionals and patients.^{185,186} The ability to laugh with a patient is a sign of good rapport. Mutual understanding while sharing some laughs may be more important than the diagnosis or formal treatment.¹⁸⁷ Laughter is also a quality-of-life and well-being enhancement therapy for both health care professionals and patients.¹⁸⁸ Laughter may help dissipate tension, fear, frustration,⁶⁴ and other stress such as “burnout,” which is becoming increasingly more common and troublesome among today’s medical staff.^{189,190} However, laughter is an often neglected resource in managing personal and professional stress. Proponents of “positive psychology” have identified humor and laughter as one of the 24 positive personal “values and attributes.”^{191,192} Therefore, health care professionals can play a significant part in eliciting better understanding of laughter benefits in clinical conditions and real-world life for both patients and themselves. But like any other skill, the effective use of laughter for therapeutic purposes needs to be learned, practiced, and developed. Practical guidelines or advice on laughter therapy have not been developed as yet to help health care professionals (and others) implement laughter techniques in their health care portfolio.

CONCLUSIONS

According to this review, the following can be concluded on the evidence-based therapeutic value of laughter in medicine:

- Current empirical data for the psychological benefits associated with laughter is stronger than that of its physiological benefits; however, further well-designed research is warranted in all of these areas.
- Overall, there are not enough research findings to conclude that laughter is an all-around healing agent, but there is sufficient evidence to suggest that laughter has some positive, quantifiable effects on certain aspects of health.
- Laughter as a medicine, as a mind-body therapy for health care, is almost never used in “traditional” clinical settings and often overlooked as a form of CAM.
- In this era of evidence-based medicine, it seems appropriate that laughter therapy takes its place as a CAM discipline in the prevention and treatment of illnesses.
- The following seems to be good advice for both patients and health care professionals: “Add laughter to your working and daily life, remember to laugh regularly, share your laughs, and help others to laugh, too.”

REFERENCES

1. Cousins N. Anatomy of an illness (as perceived by the patient). *N Engl J Med*. 1976;295(26):1458-1463.
2. Rosner F. Therapeutic efficacy of laughter in medicine. *Cancer Invest*. 2002;20(3):434-436.
3. Goodman J, Fry WF Jr. Toward optimal health: the experts discuss therapeutic humor. Interview by Jodi R. Godfrey. *J Women's Health (Larchmt)*. 2004;13(5):474-479.
4. Martin RA. Sense of humor and physical health: theoretical issues recent findings, and

- future directions. *Humor*. 2004;17(1):1-20.
5. Saper B. Humor in psychiatric healing. *Psychiatr Q*. 1988;59(4):306-319.
6. Martin RA. Humor, laughter, and physical health: methodological issues and research findings. *Psychol Bull*. 2001;127(4):504-519.
7. Provine RR. *Laughter: A Scientific Investigation*. New York, NY: Penguin Books; 2000.
8. Salovey P, Rothman AJ, Detweiler JB, Steward WT. Emotional status and physical health. *Am Psychologist*. 2000;55(1):110-121.
9. Berk R, Martin R, Baird D, Nozik B. *What Everyone Should Know About Humor and Laughter*. Aliso Viejo, CA: Association for Applied and Therapeutic Humor; 2008.
10. Provine R. *Laughter*. *Amer Scientist*. 1996 Jan-Feb;84:38-48.
11. Mahony DL, Burroughs WJ, Lippman LG. Perceived attributes of health-promoting laughter: a cross-generational comparison. *J Psychol*. 2002;136(2):171-181.
12. Mora-Ripoll R, García-Rodera MC. Therapeutic value of laughter in medicine [article in Spanish]. *Med Clin (Barc)*. 2008;131(18):694-698.
13. García-Rodera MC. *El Poder de la Risa*. Barcelona, Spain: Morales i Torres; 2003.
14. No authors listed. Reviews: making sense of an often tangled skein of evidence. *Ann Intern Med*. 2005;142(12 Pt 1):1019-1020.
15. Dixon-Woods M, Agarwal S, Jones D, Young B, Sutton A. Synthesising qualitative and quantitative evidence: a review of possible methods. *J Health Serv Res Policy*. 2005;10(1):45-53.
16. Fry WF. The biology of humor. *Humor*. 1994;7(2):111-126.
17. Sahakian A, Frishman WH. Humor and the cardiovascular system. *Altern Ther Health Med*. 2007;13(4):56-58.
18. Kataria M. *Laugh for No Reason*. Mumbai, India: Madhuri International; 2005.
19. Martin RA. *The Psychology of Humor: An Integrative Approach*. London, England: Elsevier Academic Press; 2006.
20. Zweyer K, Velker B, Ruch W. Do cheerfulness, exhilaration, and humor production moderate pain tolerance? A FACS study. *Humor*. 2004;17(1-2):85-119.
21. Cohen S. Psychosocial models of the role of social support in the etiology of physical disease. *Health Psychol*. 1988;7(3):269-297.
22. Cohen S, Underwood LG, Gottlieb BH, eds. *Social Support Measurement and Intervention: A Guide for Health and Social Scientists*. New York, NY: Oxford University Press; 2000.
23. Penson RT, Partridge RA, Rudd P, et al. Laughter: the best medicine? *Oncologist*. 2005;10(8):651-660.
24. Trent B. Ottawa lodges add humour to armamentarium in fight against cancer. *Can Med Assoc J*. 1990;142(2):163-164, 166.
25. Erdman L. Laughter therapy for patients with cancer. *Oncol Nurs Forum*. 1991;18(8):1359-1363.
26. Bellert JL. Humor. A therapeutic approach in oncology nursing. *Cancer Nurs*. 1989;12(2):65-70.
27. Kimata H. Effect of humor on allergen-induced wheal reactions. *JAMA*. 2001;285(6):738.
28. Kimata H. Laughter elevates the levels of breast-milk melatonin. *J Psychosom Res*. 2007;62(6): 699-702.
29. Bennett MP, Zeller JM, Rosenberg L, McCann J. The effect of mirthful laughter on stress and natural killer cell activity. *Altern Ther Health Med*. 2003;9(2):38-45.
30. McClelland R, Cheriff A. The immunoenhancing effects of humour on secretory IgA and resistance to respiratory infections. *Psychol Health*. 1997;12(3):329-344.
31. Bukutu C, Le C, Vohra S. Asthma: a review of complementary and alternative therapies. *Ped Review*. 2008;29(8):e44-e49.
32. Boiten FA. The effects of emotional behaviour on components of the respiratory cycle. *Biol Psychol*. 1998;49(1-2):29-51.
33. Svebak S. Respiratory patterns as predictors of laughter. *Psychophysiology*. 1975;12(1):62-65.
34. Boone T, Hansen S, Erlandson A. Cardiovascular responses to laughter: a pilot project. *Appl Nurs Res*. 2000;13(4):204-208.
35. Nahas R. Complementary and alternative medicine approaches to blood pressure reduction: an evidence-based review. *Can Fam Physician*. 2008;54(11):1529-1533.
36. Hayashi T, Urayama O, Hori M, et al. Laughter modulates prorenin receptor gene expression in patients with type 2 diabetes. *J Psychosom Res*. 2007;62(6):703-706.
37. Yoshino S, Fujimori J, Kohda M. Effects of mirthful laughter on neuroendocrine and immune systems in patients with rheumatoid arthritis. *J Rheumatol*. 1996;23(4):793-794.
38. Basmajian JV. The elixir of laughter in rehabilitation. *Arch Phys Med Rehabil*. 1998;79(12):1597.
39. Schmitt N. Patients' perception of laughter in a rehabilitation hospital. *Rehabil Nurs*. 1990;15(3):143-146.
40. Saper B. The therapeutic use of humor for psychiatric disturbances of adolescents and adults. *Psychiatr Q*. 1990;61(4):261-272.
41. Gelkopf M, Kreitler S, Sigal M. Laughter in a psychiatric ward. Somatic, emotional, social, and clinical influences on schizophrenic patients. *J Nerv Ment Dis*. 1993;181(5):283-289.
42. van Alphen PJ. Humor as diagnostic and therapeutic tool in psychiatry. Laughter is healthy [article in Dutch]. *TVZ*. 1997;107(22):674-677.
43. Seaward BL. Humor's healing potential. *Health Prog*. 1992;73(3):66-70.
44. Krack P, Kumar R, Ardouin C, et al. Mirthful laughter induced by subthalamic nucleus stimulation. *Mov Disord*. 2001;16(5):867-875.
45. Fried I, Wilson CI, MacDonald KA, Behnke EJ. Electric current stimulates laughter. *Nature*. 1998;391(6668):650.
46. Taber KH, Redden M, Hurley RA. Functional anatomy of humor: positive affect and chronic mental illness. *J Neuropsychiatry Clin Neurosci*. 2007;19(4):358-362.
47. Meyer M, Baumann S, Wildgruber D, Alter K. How the brain laughs. Comparative evi-

dence from behavioral, electrophysiological and neuroimaging studies in human and monkey. *Behav Brain Res.* 2007;182(2):245-260.

48. Kawakami K, Takai-Kawakami K, Tomonaga M, Suzuki J, Kusaka T, Okai T. Origins of smile and laughter: a preliminary study. *Early Hum Dev.* 2006;82(1):61-66.
49. Pearce JM. Some neurological aspects of laughter. *Eur Neurol.* 2004;52(3):169-171.
50. Wild B, Rodden FA, Grodd W, Ruch W. Neural correlates of laughter and humor. *Brain.* 2003;126(Pt 10):2121-2138.
51. Luschei ES, Ramig LO, Finnegan EM, Baker KK, Smith ME. Patterns of laryngeal electromyography and the activity of the respiratory system during spontaneous laughter. *J Neurophysiol.* 2006;96(1):442-450.
52. Bea JA, Marijuán PC. The informational patterns of laughter. *Entropy.* 2003;5(2):205-213.
53. Kohler KJ. "Speech-smile", "speech-laugh", "laughter" and their sequencing in dialogic interaction. *Phonetica.* 2008;65(1-2):1-18.
54. Bachorowski JA, Smoski MJ, Owren MJ. The acoustic features of human laughter. *J Acoust Soc Am.* 2001;110(3):1581-1597.
55. Bachorowski JA, Owren MF. Not all laughs are alike: voiced but not unvoiced laughter readily elicits positive affect. *Psychol Sci.* 2001;12(3):252-257.
56. Williams H. Humor and healing: therapeutic effects in geriatrics. *Gerontion.* 1986;1(3):14-17.
57. O'Brien ME. Integrative geriatrics. Combining traditional and alternative medicine. *N C Med J.* 1996;57(6):364-367.
58. Tennant KF. Laugh it off. The effect of humor on the well-being of the older adult. *J Gerontol Nurs.* 1990;16(12):11-17.
59. Evans S, Tsao JC, Zeltzer LK. Complementary and alternative medicine for acute procedural pain in children. *Altern Ther Health Med.* 2008;14(5):52-56.
60. Henderson SW, Rosario K. But seriously: clowning in children's mental health. *J Am Acad Child Adolesc Psychiatry.* 2008;47(9):983-986.
61. Reddy V, Williams E, Vaughan A. Sharing humour and laughter in autism and Down's syndrome. *Br J Psychol.* 2002;93(Pt 2):219-242.
62. Beddoe AE, Lee KA. Mind-body interventions during pregnancy. *J Obstet Gynecol Neonatal Nurs.* 2008;37(2):165-175.
63. Cantó MA, Quiles JM, Vallejo OG, et al. Evaluation of the effect of hospital clown's performance about anxiety in children subjected to surgical intervention [article in Spanish]. *Cir Pediatr.* 2008;21(4):195-198.
64. Robinson V. The purpose and function of humor in OR nursing. *Today's OR Nurse.* 1993;15(6):7-12.
65. Liechty RD. Humor and the surgeon. *Arch Surg.* 1987;122(5):519-522.
66. Morse DR. Use of humor to reduce stress and pain and enhance healing in the dental setting. *J N J Dent Assoc.* 2007;78(4):32-36.
67. Wilde JA. Laughing all the way. *Dent Today.* 1998;17(12):38-41.
68. Davis DM, Fiske J, Scott B, Radford DR. The emotional effects of tooth loss in a group of partially dentate people: a quantitative study. *Eur J Prosthodont Restor Dent.* 2001;9(2):53-57.
69. McCreaddie M, Wiggins S. The purpose and function of humour in health, health care and nursing: a narrative review. *J Adv Nurs.* 2008;61(6):584-595.
70. MacDonald CM. A chuckle a day keeps the doctor away: therapeutic humor and laughter. *J Psychosoc Nurs Ment Health Serv.* 2004;42(3):18-25.
71. Mathew FM. Laughter is the best medicine: the value of humour in current nursing practice. *Nurs J India.* 2003;94(7):146-147.
72. Astedt-Kurki P, Isola A. Humour between nurse and patient, and among staff: analysis of nurses' diaries. *J Adv Nurs.* 2001;35(3):452-458.
73. Bakerman HM. Humour as a nursing intervention. *Axone.* 1997;18(3):56-61.
74. Herth KA. Laughter. A nursing Rx. *Am J Nurs.* 1984; 84(8):991-992.
75. Berry S. Humor me on this one. The healing role of humor in EMS. *JEMS.* 2003;28(11):66-79.
76. Ratzan RM. Laughter in the temple of pain and illness. *Emerg Med.* 1996;14(1):92-95.
77. Leiber DB. Laughter and humor in critical care. *Dimens Crit Care Nurs.* 1986;5(3):162-170.
78. Borod M. SMILES—toward a better laughter life: a model for introducing humor in the palliative care setting. *J Cancer Educ.* 2006;21(1):30-34.
79. Dean RA, Gregory DM. Humor and laughter in palliative care: an ethnographic investigation. *Palliat Support Care.* 2004;2(2):139-148.
80. Dean RA. Humor and laughter in palliative care. *J Palliative Care.* 1997;13(1):34-39.
81. Herth K. Contributions of humor as perceived by the terminally ill. *Am J Hosp Care.* 1990;7:36-40.
82. Graham LL, Cates JA. Responding to the needs of the terminally ill through laughter and play. *Am J Hosp Care.* 1989;6(1):29-30.
83. Vagnoli L, Caprilli S, Robiglio A, Messeri A. Clown doctors as a treatment for preoperative anxiety in children: a randomized, prospective study. *Pediatrics.* 2005;116(4):e563-e567.
84. Parish AB. It only hurts when I don't laugh. *Am J Nurs.* 1994;94(8):46-47.
85. Strickland D. Overcome terminal seriousness: let go, laugh, and lighten up! *Sem Perioper Nurs.* 1999;8(2):53-59.
86. Hunter P. Humor therapy in home care. *Caring.* 1997;16(9):56-57.
87. Ruiz Gómez MC, Rojo Pascual C, Ferrer Pascual MA, Jiménez Navascués L, Ballesteros García M. Complementary therapies in the cares: humour and laugh therapy [article in Spanish]. *Index Enferm (Gran).* 2005;14(48-49):37-41.
88. Harries G. Use of humour in patient care. *Br J Nurs.* 1995;4(17):984-986.
89. Mallett J. Use of humour and laughter in patient care. *Br J Nurs.* 1993;2(3):172-175.
90. Granek-Catarivas M, Goldstein-Ferber S, Azuri Y, Vinker S, Kahan E. Use of humour in primary care: different perceptions among patients and physicians. *Postgrad Med J.* 2005;81(952):126-130.
91. Fowler JH, Christakis NA. Dynamic spread of happiness in a large social network: longitudinal analysis over 20 years in the Framingham Heart Study. *BMJ.* 2008 Dec 4;337:a2338.
92. Chapel M, Batten M, Brown J, et al. Frequency of public laughter in relation to sex, age, ethnicity, and social context. *Percept Mot Skills.* 2002;95(3 Pt 1):746.
93. Demmer C. A survey of complementary therapy services offered by hospices. *J Palliat Med.* 2004;7(4):510-516.
94. Astin JA, Shapiro SL, Eisenberg DM, Forsy KL. Mind-body medicine: state of the science, implications for practice. *J Am Board Fam Pract.* 2003;16(2):131-147.
95. Zagvazdin Y. Movies and emotional engagement: laughing matters in lecturing. *Fam Med.* 2007;39(4):245-247.
96. Park CM. Diversity, the individual, and proof of efficacy: complementary and alternative medicine in medical education. *Am J Public Health.* 2002;92(10):1568-1572.
97. Tiralongo E, Wallis M. Integrating complementary and alternative medicine education into the pharmacy curriculum. *Am J Pharm Edu.* 2008;72(4):74.
98. Paskind HA. Effects of laughter on muscle tone. *Arch Neurol Psychiatry.* 1932;28(3):623-628.
99. Overeem S, Taal W, Ocal Gezici E, Lammers G, Van Dijk J. Is motor inhibition during laughter due to emotional or respiratory influences? *Psychophysiology.* 2004;41(2):254-258.
100. Fry W, Savin W. Mirthful laughter and blood pressure. *Humor.* 1988;1:49-62.
101. Fry WF, Rader C. The respiratory components of mirthful laughter. *J Biol Psychol.* 1977; 19(2):39-50.
102. Fry WF Jr. The physiologic effects of humor, mirth, and laughter. *JAMA.* 1992;267(13):1857-1858.
103. Sakuragi S, Sugiyama Y, Takeuchi K. Effects of laughing and weeping on mood and heart rate variability. *J Physiol Anthropol Appl Human Sci.* 2002;21(3):159-165.
104. Filippelli M, Pellegrino R, Iandelli I, et al. Respiratory dynamics during laughter. *J Appl Physiol.* 2001;90(4):1441-1446.
105. Fry WF Jr, Stoff PE. Mirth and oxygen saturation levels of peripheral blood. *Psychother Psychosom.* 1971;19(1):76-84.
106. Yovetich NA, Dale JA, Hudak MA. Benefits of humor in reduction of threat-induced anxiety. *Psychol Rep.* 1990;66(1):51-58.
107. McMahon C, Mahmud A, Feely J. Taking blood pressure—no laughing matter! *Blood Press Monit.* 2005;10(2):109-110.
108. White S, Camarena P. Laughter as a stress reducer in small groups. *Humor. Int J Humor Res.* 1989;2(1):73-79.
109. Kajiwara M, Ohira T, Sato S, Tanaka H, Shimamoto T. Effects of laughter and physical exercise on health status among healthy elderly men and women [abstract]. *Auton Neurosci.* 2007;135(1-2):150.
110. Buchowski MS, Majchrzak KM, Blomquist K, Chen KY, Byrne DW, Bachorowski JA. Energy expenditure of laughter. *Int J Obes (Lond).* 2007;31(1):131-137.
111. Miller M, Mangano C, Park Y, Goel R, Plotnick GD, Vogel RA. Impact of cinematic viewing on endothelial function. *Heart.* 2006;92(2):261-262.
112. Xaplanteris P, Vlachopoulos C, Galifou I, et al. Beneficial effect of laughter on cortisol and von Willebrand factor levels: a possible link with endothelial function. *Eur Heart J.* 2007;28(Abtract supplement):588.
113. Xaplanteris P, Vlachopoulos C, Baou K, et al. Divergent effects of laughter and mental stress on aortic stiffness and wave reflections. *Eur Heart J.* 2007;28(Abtract supplement):591.
114. Clark A, Seidler A, Miller M. Inverse association between sense of humor and coronary heart disease. *Int J Cardiol.* 2001;80(1):87-88.
115. Tan SA, Tan LG, Berk LS. Mirthful laughter an effective adjunct in cardiac rehabilitation. *Can J Cardiol.* 1997;13(suppl B):190.
116. Berk LS, Tan L, Tan S. Mirthful laughter, as adjunct therapy in diabetic care, attenuates catecholamines, inflammatory cytokines, C-RP, and myocardial infarction occurrence [abstract]. *FASEB J.* 2008;22(Meeting abstracts):1226.2.
117. Berk LS, Felten DL, Tan SA, Bittman BB, Westengard J. Modulation of neuroimmune parameters during the eustress of humor-associated mirthful laughter. *Altern Ther Health Med.* 2001;7(2):62-72, 74-76.
118. Berk LS, Tan SA, Fry WF, et al. Neuroendocrine and stress hormone changes during mirthful laughter. *Am J Med Sci.* 1989;298(6):390-396.
119. Berk LS, Tan SA, Nehlsen-Cannarella SL, et al. Humor associated laughter decreases cortisol and increases spontaneous lymphocyte blastogenesis. *Clin Res.* 1988;36:435A.
120. Toda M, Kusakabe S, Nagasawa S, Kitamura K, Morimoto K. Effect of laughter on endocrinological stress marker chromogranin A. *Biomed Res.* 2007;28(2):115-118.
121. Levi L. The urinary output of adrenalin and noradrenalin during pleasant and unpleasant emotional states: a preliminary report. *Psychosom Med.* 1965 Jan-Feb;27:80-85.
122. Berk LS, Tan SA. [beta]-endorphin and HGH increase are associated with both the anticipation and experience of mirthful laughter [abstract]. *FASEB J.* 2006;20(4):A382.
123. Berk LS, Tan SA, Berk D. Cortisol and catecholamine stress hormone decrease is associated with the behaviour of perceptual anticipation of mirthful laughter [abstract]. *FASEB J.* 2008;22(Meeting abstracts):946.11.
124. Takahashi K, Iwase M, Yamashita K, et al. The elevation of natural killer cell activity induced by laughter in a crossover-designed study. *Int J Molecular Med.* 2001;8(6):645-650.
125. Wise B. Comparison of immune response to mirth and to distress in women at risk for recurrent breast cancer. *Diss Abstr Int.* 1989;49:2918.
126. Lefcourt H, Davidson-Katz K, Kueneman K. Humor and immune system functioning. *Humor. Int J Humor Res.* 1990;3:305-321.
127. Dillon KM, Minchoff B, Baker KH. Positive emotional status and enhancement of the immune system. *Int J Psychiatry Med.* 1985-1986;15(1):13-18.
128. Labott SM, Ahleman S, Wolever ME, Martin RB. The physiological and psychological effects of the expression and inhibition of emotion. *Behav Med.* 1990;16(4):182-189.
129. Martin RA, Dobbins JP. Sense of humor, hassles, and immunoglobulin A: evidence for a

- stress-moderating effect of humor. *Int J Psychiatry Med.* 1988;18(2):93-105.
130. Weisenberg M, Raz T, Hener T. The influence of film-induced mood on pain perception. *Pain.* 1998;76(3):365-375.
 131. Rotton J, Shats M. Effects of state humor, expectancies, and choice on postsurgical mood and self-medication: a field experiment. *J Appl Social Psychol.* 1996;26(20):1775-1794.
 132. Nevo O, Keinan G, Teshimovsky-Arditi M. Humor and pain tolerance. *Humor.* 1993;6(1):71-88.
 133. Cogan R, Cogan D, Waltz W, McCue M. Effects of laughter and relaxation on discomfort thresholds. *J Behav Med.* 1987;10(2):139-144.
 134. Van Zandt S, LaFont C. Can a laugh a day keep the doctor away? *J Pract Nurs.* 1985;35(3):32-35.
 135. Hudak DA, Dale JA, Hudak MA, DeGood DE. Effects of humorous stimuli and sense of humor on discomfort. *Psychol Rep.* 1991;69(3 Pt 1):779-786.
 136. Kimata H. Effect of viewing a humorous vs. nonhumorous film on bronchial responsiveness in patients with bronchial asthma. *Physiol Behav.* 2004;81(4):681-684.
 137. Brutsche MH, Grossman PG, Müller RE, et al. Impact of laughter on air trapping in severe chronic obstructive lung disease. *Int J COPD.* 2008;3(1):185-192.
 138. Matsuzaki T, Nakajima A, Ishigami S, Tanno M, Yoshino S. Mirthful laughter differentially affects serum pro- and anti-inflammatory cytokine levels depending on the level of disease activity in patients with rheumatoid arthritis. *Rheumatology (Oxford).* 2006;45(2):182-186.
 139. Ishigami S, Nakajima A, Tanno M, Matsuzaki T, Suzuki I, Yoshino S. Effects of mirthful laughter on growth hormone, IGF-1 and substance P in patients with rheumatoid arthritis. *Clin Exp Rheumatol.* 2005;23(5):651-657.
 140. Nakajima A, Hirai H, Yoshino S. Reassessment of mirthful laughter in rheumatoid arthritis. *J Rheumatol.* 1999;26(2):521-513.
 141. Hayashi K, Hayashi T, Iwanaga S, et al. Laughter lowered the increase in postprandial blood glucose. *Diabetes Care.* 2003;26(11):1651-1652.
 142. Hayashi T, Urayama O, Kawai K, et al. Laughter regulates gene expression in patients with type 2 diabetes. *Psychother Psychosom.* 2006;75(1):62-65.
 143. Nasir UM, Iwanaga S, Nabi AH, et al. Laughter therapy modulates the parameters of renin-angiotensin system in patients with type 2 diabetes. *Int J Mol Med.* 2005;16(6):1077-1081.
 144. Hayashi T, Tsujii S, Iburi T, et al. Laughter up-regulates the genes related to NK cell activity in diabetes. *Biomed Res.* 2007;28(6):281-285.
 145. Kimata H. Reduction of allergen-specific IgE production by laughter. *Eur J Clin Invest.* 2004;34(9):76-77.
 146. Kimata H. Viewing humorous film improves nighttime wakening in children with atopic dermatitis. *Indian Pediatr.* 2007;44(4):281-285.
 147. Kimata H. Reduction of plasma levels of neurotrophins by laughter in patients with atopic dermatitis. *Pediatr Asthma Allergy Immunol.* 2004;17(2):131-135.
 148. Marci CD, Moran EK, Orr SP. Physiologic evidence for the interpersonal role of laughter during psychotherapy. *J Nerv Ment Dis.* 2004;192(10):689-695.
 149. Pettigrew JD. Laughter abolishes binocular rivalry. *Clin Exp Optom.* 2005;88(1):39-45.
 150. Averill JR. Autonomic response patterns during sadness and mirth. *Psychophysiology.* 1969;5(4):399-414.
 151. Foley E, Matheis R, Schaefer C. Effect of forced laughter on mood. *Psychol Rep.* 2002;90(1):184.
 152. Neuhoff CC, Schaefer C. Effects of laughing, smiling, and howling on mood. *Psychol Rep.* 2002;91(3 Pt 2):1079-1080.
 153. Szabo A, Ainsworth SE, Danks PK. Experimental comparison of the psychological benefits of aerobic exercise, humor, and music. *Humor.* 2005;18(3):235-246.
 154. Morgan AJ, Jorm AF. Self-help interventions for depressive disorders and depressive symptoms: a systematic review. *Ann Gen Psychiatry.* 2008 Aug 19;7:13.
 155. Abel MH. Interaction of humor and gender in moderating relationships between stress and outcomes. *J Psychol.* 1998;132(3):267-276.
 156. Kuiper NA, Martin RA. Laughter and stress in daily life: relation to positive and negative affect. *Motiv Emot.* 1998;22(2):133-153.
 157. Moore DB. Make them laugh. Therapeutic humor for patients with grief-related stress or anxiety. *Adv Nurse Pract.* 2000;8(8):34-37.
 158. Gelkopf M, Sigal M, Kramer R. Therapeutic use of humor to improve social support in an institutionalized schizophrenic inpatient community. *J Soc Psychol.* 1994;134(2):175-182.
 159. Gelkopf M, Gonen B, Kurs R, Melamed Y, Bleich A. The effect of humorous movies on inpatients with chronic schizophrenia. *J Nerv Ment Dis.* 2006;194(11):880-883.
 160. Abel MH. Humor, stress and coping strategies. *Humor.* 2002;15(4):365-381.
 161. Nezelek JB, Derks P. Use of humor as a coping mechanism, psychological adjustment, and social interaction. *Humor.* 2001;14(4):395-413.
 162. Arminen I, Halonen M. Laughing with and at patients: the role of laughter in confrontations in addiction group therapy. *Qual Rep.* 2007;12(3):484-513.
 163. Ventis WL, Higbee G, Murdock SA. Using humor in systematic desensitization to reduce fear. *J Gen Psychol.* 2001;128(2):241-253.
 164. Walter M, Hänni B, Haug M, et al. Humour therapy in patients with late-life depression or Alzheimer's disease: a pilot study. *Int J Geriatr Psychiatry.* 2007;22(1):77-83.
 165. Takayanagi K. Laughter education and the psycho-physical effects: introduction of smile-sun method. *Jpn Hosp.* 2007 Dec;(26):31-35.
 166. Scholl JC, Ragan SL. The use of humor in promoting positive provider-patient interactions in a hospital rehabilitation unit. *Health Commun.* 2003;15(3):319-330.
 167. Amaki M, Kamide K, Takiuchi S, Niizuma S, Horio T, Kawano Y. A case of neurally mediated syncope induced by laughter successfully treated with combination of propranolol and midodrine. *Int Heart J.* 2007;48(1):123-127.
 168. Sarzi Braga S, Manni R, Pedretti RF. Laughter-induced syncope. *Lancet.* 2005;366(9483):426.
 169. Bloomfield D, Jazrawi S. Shear hilarity leading to laugh syncope in a healthy man. *JAMA.* 2005;293(23):2863-2864.
 170. Cox SV, Eisenhauer AC, Hreib K. "Seinfeld syncope." *Catheter Cardiovasc Diagn.* 1997;42(2):242.
 171. Holden R. *La Risa, la Mejor Medicina.* Barcelona, Spain: Ediciones Oniro; 1999.
 172. Temprano E. *El Arte de la Risa.* Barcelona, Spain: Editorial Seix Barral; 1999.
 173. García Larrauri B. *Programa para Mejorar el Sentido del Humor.* Madrid, Spain: Ediciones Pirámide; 2008.
 174. Ekman P, Davidson RJ, Friesen WV. The Duchenne Smile: emotional expression and brain physiology II. *J Pers Soc Psychol.* 1990;58(2):342-353.
 175. García-Rodera MC. *Cómo Beneficiarse del Poder de la Risa.* Barcelona, Spain: Morales i Torres Editores; 2007.
 176. Wortzel HS, Oster TJ, Anderson CA, Arciniegas DB. Pathological laughing and crying: epidemiology, pathophysiology and treatment. *CNS Drugs.* 2008;22(7):531-545.
 177. Gendry S. *Laughter Yoga 101: From 0 to Laughter in 0.5 Seconds.* Los Angeles, CA: American School of Laughter Yoga; 2007.
 178. Solomon G. Psychoneuroimmunology: interactions between the central nervous system and immune system. *J Neurosci Res.* 1987;18(1):1-9.
 179. Martin RA. Humor and health. In: Raskin V, ed. *The Primer of Humor Research.* New York, NY: Mouton de Gruyter; 2008:479-521.
 180. Beckman H, Regier N, Young J. Effect of workplace laughter groups on personal efficacy beliefs. *J Prim Prev.* 2007;28(2):167-182.
 181. Nagendra HR, Chaya MS, Nagarathna R, Kataria M, Manjunath MK, Rao R. *The Efficacy of Laughter Yoga on IT Professionals to Overcome Stress.* Mumbai, India: Laughter Yoga International; 2007.
 182. Kimata H, Morita A, Furuhashi S, Itakura H, Ikenobu K, Kimura K. Assessment of laughter by diaphragm electromyogram. *Eur J Clin Invest.* 2009;39(1):78-79.
 183. Balick MJ, Lee R. The role of laughter in traditional medicine and its relevance to the clinical setting: healing with ha! *Altern Ther Health Med.* 2003;9(4):88-91.
 184. Mora-Ripoll R. Laughter medicine and therapy: A complementary/alternative approach in health and patient care [article in Spanish]. *Med Clin (Barc).* 2009 Sep 21. [Epub ahead of print]
 185. Sanz Ortiz J. Therapeutic value of humor [article in Spanish]. *Med Clin (Barc).* 2002;119(19):734-737.
 186. Karl KA, Peluchette JV, Harland L. Is fun for everyone? Personality differences in health-care providers' attitudes toward fun. *J Health Hum Serv Adm.* 2007;29(4):409-447.
 187. Elliott-Binns CP. Laughter and medicine. *J R Coll Gen Pract.* 1985;35(277):364-365.
 188. Richman J. The lifesaving function of humor with the depressed and suicidal elderly. *Gerontologist.* 1995;35(2):271-273.
 189. Esteve M, Larraz C, Soler J, Yaman H. Burn-out in Spanish general practitioners [article in Spanish]. *Aten Primaria.* 2005;35(2):108-109.
 190. Casado Moragón A, Muñoz Durán G, Ortega González N, Castellanos Asenjo A. Burnout in professionals belonging to palliative, intensive care and emergency services in a general hospital [article in Spanish]. *Med Clin (Barc).* 2005;16:124(14):554-555.
 191. de Vera Poseck B. *Psicología Positiva: Una Nueva Forma de Entender la Psicología.* Madrid, Spain: Calamar Ediciones; 2008.
 192. Seligman ME, Steen RA, Park N, Peterson C. Positive psychology progress: empirical validation of interventions. *Am Psychol.* 2005;60(5):410-421.